## Dr Jack D. Singer LTD

MD FRCPCH
Consultant Paediatrician
Practice Associate: Dr Olga Kapellou

(020) 7937 9406 (home)

73 Harley Street London W1G 8QJ Tel: (020) 7935 2023 Fax: (020) 7935 3857

## Meningitis B Vaccination

We are now able to offer the Meningitis B vaccination for your child/children. The Meningococcal B (MenB) infection is the leading cause of bacterial meningitis and septicaemia in the UK, and this kills more children under 5 than any other infectious diseases in the UK.

Bexsero Meningococcal Group B vaccine is now available and protects against invasive meningitis disease.

The vaccine is indicated for active immunisation of individuals from 2 months of age and older against invasive meningococcal disease caused by Neisseria Meningitidis group B

Age Group	Primary Immunisation	Intervals between Primary dose	Booster
Infants, 2 months to 5 months	Three doses each of 0.5ml, with the first dose given at 2, 4 & 6months of age	Not less than 1 month	Yes, one dose between 12 and 23 months
Unvaccinated infants 6 months to 11 months	Two doses each of 0.5ml	Not less than 2 months	Yes, one dose in the second year of life with an interval of at least 2 months between the primary series and booster dose
Unvaccinated children 12 months to 23 months	Two dose each of 0.5ml	Not less than 2 months	Yes, one dose with an interval of 12 months to 23 months between primary series and booster dose
Children, 2 years to 10 years	Two dose each of 0.5ml	Not less than 2 months	Need not established
Adolescents (from 11 years of age) and adults	Two dose each of 0.5ml	Not less than 1 month	Need not established  (table from Novartis vaccines
			information leaflet)

If you are interested in having your child/children vaccinated, or would like to know more about the vaccine please contact our office on 020 7935 2023 and we will be happy to discuss this with you.

Administrative Team
Jack Donald Singer MD FRCPCH
Consultant Paediatrician